



| |
|--|
| FIFRA 2(ee) Recommendation |
| EPA Reg. No. 71512-37-74530 |
| FOR DISTRIBUTION AND USE ONLY IN: TEXAS |
| THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA. |

This recommendation for Reviton herbicide tankmixed with Dicamba is valid until December 31, 2028, or until withdrawn, cancelled, or suspended.

FOR POST-HARVEST DESTRUCTION/REMOVAL OF ENLIST COTTON STALK KILL AFTER STALK MOWING

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

| CROP | PEST | REVITON RATE OF APPLICATION | DICAMBA RATE OF APPLICATION |
|---------------|-----------------------|-----------------------------|-----------------------------|
| Enlist Cotton | Regrowth After Mowing | 2.0 – 3.0 oz/A | 1.0 lb ai of Dicamba Acid |

DIRECTION FOR USE:

- Apply tank-mix of Reviton + Dicamba to Enlist cotton regrowth after stalk mowing.
- Apply to regrowth when it reaches 3 – 4 inches tall.
- **Always** include a high-quality MSO at 1 pt/A.
- Ground application only in a minimum of 15 GPA.

RESTRICTIONS:

- Two applications may be needed for effective control at a two to three-week interval.
- Do not apply Reviton at a rate greater than 3 fl oz per acre. Do not exceed 9 fl oz per acre per year.
 - Do not apply more than 2 applications of dicamba per year.
 - Do not exceed a combined total of 1.0 pound of dicamba a.e. per acre per application. Do not exceed a total of 2.0 pounds of dicamba a. e. per acre per year.
 - Do not allow livestock to graze for 21 days after application.

A tank-mix of Reviton + Dicamba + an MSO adjuvant will effectively control Enlist cotton regrowth after stalk cutting. Oftentimes the cut stalk will start regrowing again after the first application of the mix listed above and a second application will be required approximately 14 – 28 days after the first application. Thus, for best control a two-application program is recommended for most effective control.

USER OF TANKMIX MUST FOLLOW THE LABEL WITH THE MOST RESTRICTIVE USE REQUIREMENTS.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED REVITON AND DICAMBA LABELS MUST BE FOLLOWED.

THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

HELM Agro US, Inc.
401 E. Jackson St., Suite 1400
Tampa, FL 33602
Phone: 813.621.8846
Fax: 813.621.0763
info@helmagro.com
helmagro.com

Revised: 1/7/2025