



# Reviton

Supplemental Labeling

## SUPPLEMENTAL DIRECTIONS FOR THE USE OF REVITON (EPA Reg. No. 71512-37-74530)

### FOR APPLICATION ON PULSES, DRIED SHELLED PEAS FOR CONTROL OF WEEDS LISTED ON ATTACHED PAGES

### KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCIÓN

This Supplemental Label Expires on Dec 31st, 2028 and must not be used or distributed after this date.

#### Directions for Use:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of pesticide application.

Read the label affixed to the container for Reviton before applying.

Use of Reviton according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Reviton.

#### Pulses, Dried Shelled Peas

Pea (*Pisum* spp.; dry pea, field pea); chickpea (garbanzo); and lentil

Application Timing	Rate Range (fl oz/A)	Additional Information & Restrictions
Preplant through preemergence	1.0 to 2.0	<ul style="list-style-type: none"><li>• Apply as a broadcast spray using conventional low-pressure ground spray equipment or by aerial application.</li><li>• Follow manufacturer's recommendations for spraying pressure.</li><li>• Review and follow restrictions from the spray drift management section.</li><li>• The use of 2.0 fl oz requires a minimum of a 7 day preplant interval.</li><li>• Do not reapply within 14 days.</li><li>• Use higher rate for dense and/or mature weed infestations.</li><li>• Do not exceed 2.0 fl oz of product (0.044 lb ai) per acre per application.</li><li>• Do not exceed 4.0 fl oz of product (0.088 lb ai) per acre per year applied preplant through preemergence.</li></ul>
Harvest Aid / Desiccation / Defoliation	1.0 to 2.0	<ul style="list-style-type: none"><li>• Apply a broadcast spray using conventional low-pressure ground spray equipment or by aerial application.</li><li>• Follow manufacturer's recommendations for spraying pressure.</li><li>• Review and follow restrictions from the spray drift management section.</li></ul>

		<ul style="list-style-type: none"> <li>• Do not apply within 3 days of harvest.</li> <li>• Do not reapply within 7 days.</li> <li>• Inadequate coverage of foliage will result in unacceptable crop desiccation/defoliation.</li> <li>• Do not allow desiccation-treated pea vines to be grazed or fed to livestock.</li> <li>• Do not apply aid/desiccation to green lentil varieties.</li> <li>• Do not exceed 2.0 fl oz of product (0.044 lb ai) per acre per application.</li> <li>• Do not exceed a total of 2.0 fl oz of product (0.044 lb ai) per acre per year applied as harvest aid/desiccation/defoliation.</li> </ul>
<b>Harvest Aid / Desiccation / Defoliation:</b> Spray over the top of crops that have reached physiological maturity (peas with at least 80% yellow/brown pods (pod rattle stage) and no more than 30% of leaves still green, or according to Extension Service recommendations in the use area). Allow up to 10 days for optimum desiccation effect. Actual time to harvest depends on environmental and atmospheric conditions which may increase or decrease the time period stated here.		
<b>Tank Mixes:</b> Reviton is generally compatible with fertilizers and micronutrient products, provided sufficient free water is available for dispersion of all the tank mixture products. Use tank mixture combinations only when applicator experience indicates that the tank mixture will not result in objectionable crop injury.		
Refer to Tank Mixtures section for additional information regarding tank mixes.		
<b>Reviton plus Glyphosate:</b> Reviton can be applied at 1.0 to 3.0 fl oz per acre (0.023 to 0.067 lb ai per acre) in combination with glyphosate to improve overall weed control and broaden weed control spectrum. Follow glyphosate label rate and use directions (or follow local extension recommendations).		
<b>Restrictions:</b> DO NOT apply this product to residential areas. DO NOT apply this product through any type of irrigation system.		
<p>A 50-foot buffer for ground applications and a 150-foot buffer for aerial applications must be maintained between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (including grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas, shrub lands, croplands), semi-aquatic, and estuarine/marine habitats.</p> <p>To prevent potential for run-off, maintain a 25-foot vegetative filter strip between the treated field and any sensitive aquatic habitat such as but not limited to; lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries and commercial fish ponds.</p> <p>In the State of New York: Aerial applications are prohibited. Maintain a 100-foot buffer for ground applications between the point of direct application and the closest edge of sensitive aquatic habitats such as but not limited to; lakes reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries and commercial fish ponds.</p>		

Warranty and Limitation of Damages: Seller warrants to those persons lawfully acquiring title to this product that at the time of first sale of this product by Seller that this product conformed to its chemical description and was reasonably fit for the purposes stated on the label when used in accordance with Seller's directions under normal conditions of use. To the extent consistent with applicable law, Buyers and users of this product assume the risk of any use contrary to such directions. EXCEPT AS PROVIDED ELSEWHERE IN WRITING CONTAINING AN EXPRESS REFERENCE TO THIS WARRANTY AND LIMITATION OF DAMAGES, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OR GUARANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY, AND NO AGENT OF SELLER IS AUTHORIZED TO DO SO. In no event shall Seller's liability for any breach of warranty or guaranty exceed the purchase price of the product as to which a claim is made. To the extent consistent with applicable law, Buyers and users of this product are responsible for all loss or damage from use or handling of this product which results from conditions beyond the control of Seller, including, but not limited to, incompatibility with other products unless otherwise expressly provided in Directions for Use of this product, weather conditions, cultural practices, moisture conditions or other environmental conditions outside of the ranges that are generally recognized as being conducive to good agricultural and/or horticultural practices.

## WEED LIST

Broadleaf and grass weeds controlled (C) or suppressed (S) by applications of Reviton applied to actively growing weeds at 2.0 to 3.0 fl oz per acre or at 1.0 to 3.0 fl oz per acre in tank mix with glyphosate.

	Common Name	Scientific Name	Reviton	Reviton+ glyphosate <sup>1</sup>
<b>Broadleaf Weeds</b>	Amaranth, Palmer	<i>Amaranthus palmeri</i>	S-C <sup>2</sup>	S-C <sup>3</sup>
	Buttercup spp	<i>Ranunculus spp</i>	S	C
	Canola, volunteer	<i>Brassica rapa</i>	C	C
	Chickweed, common	<i>Stellaria media</i>	S	C
	Chickweed, mouse-ear	<i>Cerastium fontanum</i>	S	C
	Clover, white	<i>Trifolium repens</i>	S-C	C
	Dandelion, common	<i>Taraxacum officinale</i>	S-C	C
	Deadnettle, purple	<i>Lamium purpureum</i>	C	C
	Dock, curly	<i>Rumex crispus</i>	S	S-C
	Eveningprimrose, cutleaf	<i>Oenothera laciniata</i>	S-C	C
	Geranium, Carolina	<i>Geranium carolinianum</i>	S	C
	Groundsel, cressleaf	<i>Packera glabella</i>	C	C
	Henbit, common	<i>Lamium amplexicaule</i>	C	C
	Horseweed (maretail) <sup>4</sup>	<i>Erigeron canadensis</i>	S	S-C
	Kochia	<i>Bassia scoparia</i>	S	S-C <sup>5</sup>
	Lambsquarters, common	<i>Chenopodium album</i>	C	C
	Morningglory spp	<i>Ipomoea spp</i>	C	C
	Pennycress, field	<i>Thlaspi arvense</i>	C	C
	Pigweed, redroot	<i>Amaranthus retroflexus</i>	S-C	C
	Prickly sida (teaweed)	<i>Sida spinosa</i>	C	C
	Purslane, common	<i>Portulaca oleracea</i>	S-C	C
	Radish, wild	<i>Raphanus raphanistrum</i>	S	S-C
	Ragweed, common	<i>Ambrosia artemisiifolia</i>	C <sup>2</sup>	C <sup>3</sup>
	Ragweed, giant	<i>Ambrosia trifida</i>	C	C
	Sesbania, hemp	<i>Sesbania herbacea</i>	C	C
	Shepherd's-purse	<i>Capsella bursa-pastoris</i>	C	C
	Swinecress	<i>Lepidium spp</i>	S-C	C
	Thistle, Russian	<i>Salsola kali</i>	S-C	S-C
	Velvetleaf	<i>Abutilon theophrasti</i>	C	C
	Vetch	<i>Vicia spp.</i>	S-C	C
	Waterhemp	<i>Amaranthus tuberculatus</i>	S-C <sup>2</sup>	S-C <sup>3</sup>
<b>Grass Weeds</b>	Barley, little	<i>Hordeum pusillum</i>	S	C
	Barnyardgrass	<i>Echinochloa crus-galli</i>	S	C
	Bluegrass, annual	<i>Poa annua</i>	S-C	C
	Corn, volunteer <sup>6</sup>	<i>Zea mays</i>	C	C
	Crabgrass spp.	<i>Digitaria spp.</i>	S	C
	Foxtail, giant	<i>Setaria faberi</i>	S	C
	Foxtail, green	<i>Setaria viridis</i>	S	C
	Foxtail, yellow	<i>Setaria pumila</i>	S	C
	Goosegrass	<i>Eleusine indica</i>	S	C
	Johnsongrass, seedling	<i>Sorghum halepense</i>	S	C

	Oats, wild	<i>Avena fatua</i>	C	C
	Rye, volunteer	<i>Secale cereale</i>	S	S-C
	Shattercane	<i>Sorghum bicolor</i>	S	C
	Wheat, volunteer	<i>Triticum aestivum</i>	S	C

<sup>1</sup> Refer to tank mix section for details. Rating based on glyphosate- susceptible populations.

<sup>2</sup> Except on PPO resistant populations.

<sup>3</sup> Except on glyphosate and PPO resistant populations.

<sup>4</sup> For enhanced performance on horseweed (including glyphosate resistant populations), Reviton can be mixed with 2,4-D or dicamba based products (refer to specific labels for directions/restrictions). Tank mixtures including metribuzin have also shown to be beneficial (refer to specific labels for directions/restrictions).

<sup>5</sup> For best results, target applications before Kochia has begun developing branches.

<sup>6</sup> Target application between VE-V6 stages. Add low rates of clethodim (refer to clethodim label for directions/restrictions) for more complete control. When growing point is below the ground, the addition of low rates of clethodim is required to keep regrowth from occurring.

(RV051525)

Manufactured for:  
**HELM Agro US, Inc.**  
401 E. Jackson St., Suite 1600  
Tampa, FL 33602